

TURF WEED CONTROL (continued)						
Use Stage and Herbicide	Turfgrasses	WSSA Group	Broadcast Rate/Acre		REI	Remarks and Precautions
			Amount of Formulation	Pounds Active Ingredient		
Preemergence Herbicides – Golf Course Putting Greens - continued						
fenarimol (Rubigan) 1AS (Patchwork) 0.78G	overseeded bermudagrass	n/a	See Label See Label	See Label See Label	12	Fenarimol is a systemic fungicide that reduces the infestation level of annual bluegrass. May be applied as a sequential treatment to bermudagrass greens that are overseeded with perennial ryegrass, <i>Poa Trivialis</i> or bentgrass. Sequential treatments should be applied at intervals of 10 to 14 days. Use the triple application program if bentgrass or <i>Poa Trivialis</i> is used for overseeding. For double applications, apply 6.0 fl. oz. of Rubigan AS or 6.0 lbs. of Patchwork per 1,000 sq. ft. with the last application 2 weeks prior to overseeding ryegrass. For triple applications, apply 4.0 fl. oz. of Rubigan AS or 4.0 lbs. Patchwork per 1,000 sq. ft. with the last application 2 weeks prior to overseeding ryegrass or 4 weeks prior to overseeding bentgrass or <i>Poa Trivialis</i> . A single application of Rubigan AS at 2.0 fl. oz. Or Patchwork at 2.0 lbs. per 1,000 sq. ft. in January or February will help to maintain annual bluegrass control. Rubigan will not control perennial bluegrass.
bensulide + oxadiazon (Goosegrass/Crabgrass Control) 5.25% + 1.31% G	bentgrass, bermudagrass	8+14	115.0 lbs.	6.0 bensulide + 1.5 oxadiazon	See Label	Controls summer annual grasses. Apply a light irrigation after treatment. DO NOT use on newly sprigged grasses until well established. Delay reseeding for 5 months after treatment. On putting greens overseeded with perennial ryegrass or <i>Poa Trivialis</i> apply 1/2 the maximum labeled rate to dry turf. Apply an additional application at 1/2 the maximum labeled rate to dry turf 10 to 14 days later. See label before use on putting greens.
Postemergence Herbicides – Golf Course Putting Greens						
carfentrazone (QuickSilver) 1.9 lbs./gal. (QuickSilver) 1.03 lbs./gal. (water-soluble 5 ml bag)	bentgrass	14	6.7 fl. oz. use 3-4 5 ml bags/2,000 sq. ft	0.1	Until Dry	Apply to creeping bentgrass putting greens and tees for the control of silvery-thread moss (<i>Byrum argenteum</i>). Repeat the application in 2 weeks. Add a nonionic surfactant at 0.25% v/v. Apply at a spray volume of ≥ 100 GPA. Transitory injury (yellowing) may occur when applied to bentgrass stressed from high air temperatures (> 90°F.), disease and soil moisture stress.
diclofop-methyl (Illoxan) 3EC	bermudagrass	1	1.0-1.4 qts.	0.75-1.0	See Label	Illoxan may be used for goosegrass control on bermudagrass golf greens, tees and fairways. Use the high rate for goosegrass with one to two tillers. Lower rates may be used on smaller goosegrass. DO NOT mow for 24-36 hours after application. Control requires 2-3 weeks. May cause temporary phytotoxicity (7-10 days). DO NOT tank mix with other pesticides or fertilizers as control will be reduced. DO NOT overseed treated areas for at least 2 months after the last application.
foramsulfuron (Revolver) 0.19 lb./gal.	bermudagrass	2	8.8-26.2 fl. oz.	0.013-0.04	12	Apply Revolver for: a) control of emerged annual bluegrass 2 weeks before overseeding bermudagrass with perennial ryegrass, b) postemergence control of goosegrass in late spring and early summer, c) postemergence control of annual bluegrass in non-overseeded putting greens, and d) spring transition of bermudagrass overseeded the previous fall with cool-season turfgrasses. For tillered goosegrass, make 2 applications at the high rate 7-14 days apart. DO NOT apply Revolver within 2 weeks of sprigging. Revolver is rainfast within 2 hours of application. Revolver must be allowed to dry on bermudagrass foliage before traffic is allowed through treated areas. Revolver is NOT recommended for use on creeping bentgrass greens.

Supplemental 2(ee) label recommendations may be available for some listed pesticides. It is the end-user's responsibility to consult the manufacturer or websites like www.CDMS.net for additional recommendations.

TURF WEED CONTROL (continued)						
Use Stage and Herbicide	Turfgrasses	WSSA Group	Broadcast Rate/Acre		REI	Remarks and Precautions
			Amount of Formulation	Pounds Active Ingredient		
Postemergence Herbicides – Golf Course Putting Greens - continued						
mecoprop (MCP-4 Amine)	bentgrass, bermudagrass	4	2.5 pts.	1.25	See Label	Apply to small and actively growing broadleaf weeds as a summer or winter treatment. Avoid spray drift to ornamentals tree, shrub or flower foliage. DO NOT water for 24 hours after application. Delay mowing 2-3 days before or after application. DO NOT apply during conditions of drought, high temperatures (> 90°F.) or very wet weather. Weekly applications at one-half recommended rates will provide better control of knotweed than a single application at normal use rates. DO NOT reseed for 3-4 weeks after application.
rimsulfuron (TranXit GTA) 25DF	bermudagrass	2	1.0-2.0 oz.	0.016-0.032	4	May be used: a) to control annual bluegrass before overseeding bermudagrass with perennial ryegrass or <i>Poa Trivialis</i> ; b) to control annual bluegrass in non-overseeded bermudagrass, and c) to remove perennial ryegrass or <i>Poa Trivialis</i> from bermudagrass greens in the late spring. Add a nonionic surfactant at 0.25% v/v. Apply 10-14 days in late summer or early fall before overseeding bermudagrass. DO NOT apply after overseeding, or to bentgrass putting greens. To hasten spring transition on overseeded bermudagrass, apply at the 60-75% green-up growth stage of bermudagrass, or approximately 2-3 weeks before transition is desired. Irrigate within 2 hours of a TranXit application and then follow normal irrigation schedule. TranXit is NOT recommended for use on creeping bentgrass putting greens.
MCP + 2,4-D + dicamba (Trimec Bentgrass)	bentgrass	4+4+4	2.7 pts.	0.5 + 0.15 + 0.06	See Label	Apply to small and actively-growing broadleaf weeds when air temperatures are less than 80°F. DO NOT water for 24 hours after application. DO NOT apply if bentgrass is showing the effects of environmental stresses.
sulfosulfuron (Certainty) 75WDG	bermudagrass, centipedegrass, seashore paspalum, St. Augustinegrass, zoysiagrass	2	0.75-2oz.	0.04-0.09	12	Sulfosulfuron controls nutsedge(s), annual sedges, kyllinga species, tall fescue, and certain broadleaf weeds in warm-season turfgrasses. Add a nonionic surfactant at 0.25% v/v to the spray mix. Avoid mowing for 1-2 days before and after application. For nutsedge, repeat the application at 4-10 weeks if regrowth is observed. Sulfosulfuron may be applied 7-10 days before overseeding bermudagrass with perennial ryegrass. DO NOT use on putting greens. DO NOT use on tall fescue or other cool-season grasses.
trifloxysulfuron-sodium (Monument) 75DG	bermudagrass	2	0.33-0.56 oz.	0.015-0.026	12	Controls nutsedge(s), annual sedges, kyllinga species, and certain broadleaf weeds in established bermudagrass putting greens. Applications should be made to actively-growing bermudagrass after 100% spring green-up. Monument is not recommended for use on creeping bentgrass. Add a nonionic surfactant at 0.25-0.5% v/v to the spray mix. Monument at rates of 0.1-0.3 oz./acre may be used as a spring transition aid for the removal of perennial ryegrass and <i>Poa trivialis</i> . Avoid mowing for 1-2 days before and after application. For nutsedge repeat the application at 4-6 weeks if regrowth is observed. DO NOT overseed bermudagrass with cool-season turfgrasses for 3 weeks after application.

Supplemental 2(ee) label recommendations may be available for some listed pesticides. It is the end-user's responsibility to consult the manufacturer or websites like www.CDMS.net for additional recommendations.